arXiv:1607.08555v1 [math.DG] 28 Jul 2016

ARONSON-BÉNILAN ESTIMATES FOR THE FAST DIFFUSION EQUATION UNDER THE RICCI FLOW

HUAI-DONG CAO AND MENG ZHU

Abstract. We study the fast diffusion equation with a linear forcing term,

$$\frac{\partial u}{\partial t} = \mathrm{div}(|u|^{p-1}\nabla u) + Ru, \tag{0.1}$$

under the Ricci flow on complete manifold M with bounded curvature and nonnegative curvature operator, where $0 < p < 1$ and $R = R(x,t)$ is the evolving scalar curvature of M at time t. We prove Aronson-Bénilan and Li-Yau-Hamilton type differential Harnack estimates for positive solutions of (0.1). In addition, we use similar method to prove certain Li-Yau-Hamilton estimates for the heat equation and conjugate heat equation which extend those obtained by X. Cao and R. Hamilton [10], X. Cao [9], and S. Kuang and Q. Zhang [23] to noncompact setting.

1. Introduction

This is a sequel to our paper [7], in which we derived Aronson-Bénilan and Li-Yau-Hamilton type differential Harnack estimates for positive solutions of the porous medium equation coupled with the Ricci flow on complete manifolds with bounded curvature and nonnegative curvature operator. In this paper, we study similar problems for the fast diffusion equation under the Ricci flow.

On a Riemannian manifold (M^n, g_{ij}), the fast diffusion equation (FDE) is the nonlinear parabolic equation

$$\frac{\partial u}{\partial t} = \mathrm{div}(|u|^{p-1}\nabla u), \tag{1.1}$$

where $p < 1$. Equation (1.1) is also known as the porous medium equation (PME) when $p > 1$. PME and FDE appear naturally as nonlinear diffusion models in many areas of mathematics, physics, chemistry, and engineering. For example, the PME is the model of the flow of gas through porous medium, ground water filtration, and heat radiation in plasmas, etc (see e.g. [33]), while FDE arises in plasma physics ([29]), diffusion of impurities in silicon ([22]), gas kinetics theory ([25]), etc. Moreover, in geometry, the FDE with $n = 2$ and $p = 0$ describes the Ricci flow on surfaces ([34], [17]), and it corresponds to the Yamabe flow when $n \geq 3$ and $p = \frac{n-2}{n+2}$ ([32]).

While both PME and FDE look quite similar to the heat equation, the case when $p = 1$ in (1.1), the nonlinearity and degeneracy make their existence and regularity theory differ a lot from the heat equation. On Euclidean spaces, a relatively satisfying theory for the PME has been developed over the past several decades. Readers may refer to [33] and the references therein for a thorough account of the PME. Regarding the FDE on $\mathbb{R}^n$, see e.g. [1], [21], [12], [15], [16], [13], [14], [3], and [32]. While many properties and techniques in the PME case can be shared in the FDE case, the two equations have some fundamental differences. For

The research of the first author was partially supported by NSF Grant DMS-0909581.

The research of the second author was partially supported by the National Natural Science Foundation of China Grant No.11501206.

instance, long time solution to the Cauchy problem of FDE with $0 < p < 1$ and L^1_{loc} initial condition exists without any growth restriction on the initial value ([21]), solutions of FDE with $p < 0$ and L^1 initial data (see e.g. [31]) do not exist, and the L^1 to L^∞ smoothing effect can only be carried from PME to FDE for $\frac{n-2}{n} < p < 1$ (see e.g. [32]).

If one concentrates on nonnegative solutions of (1.1) with $p > 0$, it is equivalent to the following equation:

$$\frac{\partial u}{\partial t} = \Delta u^p. \tag{1.2}$$

In the study of Cauchy problems of PME and FDE on $\mathbb{R}^n$, an essential tool is the following second order differential inequality discovered by Aronson and Bénilan ([1]):

$$\sum_i \frac{\partial}{\partial x_i}(pu^{p-2}\frac{\partial u}{\partial x_i}) \geq -\frac{\kappa}{t}, \tag{1.3}$$

where $\kappa = \frac{n}{2+n(p-1)}$, for any positive solution u to (1.2) with $p > (1-\frac{2}{n})^+$.

On a general complete Riemannian manifold (M^n, g_{ij}) with Ricci curvature bounded from below, $Rc \geq -K$ for some $K \geq 0$, such kind of differential estimate was first found by Li and Yau ([24]). Subsequently, the matrix differential Harnack estimate for the heat equation was proved by Hamilton ([20]) on Riemannian manifolds, and by Ni and the first author ([6]) on Kähler manifolds.

Unlike the heat equation, PME and FDE on Riemannian manifolds were only investigated recently. Aronson-Bénilan type estimates were first proved by Vázquez ([33]) on manifolds with nonnegative Ricci curvature, and generalized by Lu, Ni, Vázquez and Villani ([26]) to manifolds with Ricci curvature bounded from below. In particular, they showed that, if $Rc \geq -(n-1)K$ on M for some $K \geq 0$, then for any $\alpha \in (0,1)$, $\epsilon > 0$ and bounded smooth positive solution u to FDE, one has

$$\alpha\frac{v_t}{v} - \frac{|\nabla v|^2}{v} \leq \frac{\kappa\alpha^2}{C}\left[\frac{1}{t} + \frac{(1-p)(n-1)}{(1-\alpha)\epsilon}\sqrt{C}K\bar{v}_{max}\right], \tag{1.4}$$

where $v = \frac{p}{p-1}u^{p-1}$, $\kappa = \frac{n(1-p)}{2-n(1-p)}$, $\bar{v}_{max} = \max_{M\times[0,T]} v$, and $C = C(\kappa, \alpha, \epsilon) > 0$.

The study of differential Harnack estimates for parabolic equations under the Ricci flow

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} \tag{1.5}$$

on complete manifolds was originated from Hamilton's work ([18]). The readers may refer to e.g. [5], [11], [19], [27], [28], [30] for more results on differential Harnack estimates for linear and semi-linear parabolic equations under the Ricci flow and the Kähler-Ricci flow.

In recent years, among the parabolic equations coupled with the Ricci flow, the heat equation

$$\frac{\partial u}{\partial t} = \Delta u, \tag{1.6}$$

and the conjugate heat equation

$$\frac{\partial u}{\partial t} = -\Delta u + Ru \tag{1.7}$$

are two of the most studied.

In [30], Perelman first obtained a Li-Yau type estimate for the fundamental solution of the conjugate heat equation (1.7) (see also [27]). On compact manifolds, X. Cao [9] and Kuang-Zhang [23] independently showed the following Li-Yau estimate,

$$2\Delta v - |\nabla v|^2 + R \leq \frac{n}{2\tau}, \tag{1.8}$$

where $u = (4\pi\tau)^{-n/2}e^{-v}$, for arbitrary positive solutions u of (1.7).

Regarding the heat equation (1.6) under the Ricci flow, Li-Yau type estimates were first studied by Bailesteanu-X.Cao-Pulemotov [2] assuming the Ricci curvature being uniformly bounded. Most recently, Q. Zhang and the second author [35] were able to show a Li-Yau type estimate only under the assumption that the scalar curvature is uniformly bounded.

Moreover, By using Hamilton's trace differential Harnack estimate [19], X. Cao and Hamilton [10] proved the Li-Yau-Hamilton type estimate

$$2\Delta v - |\nabla v|^2 - 3R \leq \frac{2n}{t}, \tag{1.9}$$

where $v = -\ln u$ and u is any positive solution to the heat type equation

$$\frac{\partial u}{\partial t} = \Delta u + Ru \tag{1.10}$$

under the Ricci flow on a compact manifold with nonnegative curvature operator.

In [7], we studied PME (1.2) with a linear forcing term,

$$\frac{\partial u}{\partial t} = \Delta u^p + Ru, \tag{1.11}$$

where $p > 1$, under the Ricci flow on a complete manifold M^n with bounded curvature and nonnegative curvature operator and proved Aronson-Bénilan and Li-Yau-Hamilton type differential Harnack estimates for any bounded positive solution $u(x,t)$. Especially, we showed

$$\frac{|\nabla v|^2}{v} - 2\frac{v_t}{v} - \frac{R}{v} - \frac{d}{t} \leq 0,$$

where $v = \frac{p}{p-1}u^{p-1}$, $d = \max\{2\alpha, 1\}$ and $\alpha = \frac{n(p-1)}{1+n(p-1)}$. The reason for adding the term Ru in (1.11) is that the total mass $\int_M u dV$ would stay unchanged under (1.11) coupled with the Ricci flow, as it does under (1.2) in the fixed metric case.

In this paper, we consider the equation (1.11) with $0 < p < 1$, i.e., the FDE case. We prove that certain Aronson-Bénilan and Li-Yau-Hamilton type estimates as in [7] still hold. In particular, we have

Theorem 1.1. *Let $(M^n, g_{ij}(t))$, $t \in [0,T]$, be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator. If u is a bounded smooth positive solution to (1.11) with $p \in (1-\frac{1}{n}, 1)$, then for $v = \frac{p}{p-1}u^{p-1}$, we have*

$$\frac{|\nabla v|^2}{v} - 2\frac{v_t}{v} - \frac{R}{v} + KR_{max} + \frac{2\alpha}{t} \geq 0$$

on $M \times (0,T]$, where $\alpha = \frac{n(1-p)}{1-n(1-p)}$, $R_{max} = \sup_{M\times[0,T]} R$, and $K = (1-p)\max\{(\alpha+\frac{1}{2}), 2\alpha\}$.

A more general version of the differential Harnack inequality depending on a positive parameter $b > 1$ (see Theorem 2.6 and Theorem 2.11) is given in Section 2, and Theorem 1.1 is the special case when $b = 2$. It happens that the proof of these differential Harnack inequalities in the FDE case is more subtle than in the PME case, in the sense that some terms on the right hand side of Eq. (2.2) in Proposition 2.1 become bad terms in the FDE case, while they are good terms in the PME case. Thus, more effort was needed in order to have them controlled.

Remark 1.2. *When the metric is fixed, Lu-Ni-Vázquez-Villani* [26] *obtained a series of Aronson-Bénilan estimates for FDE* (1.1) *in which the parameters $b \in (0,1)$ and $p \in (1-\frac{2}{n}, 1)$ (see* (1.4)*). However, under the Ricci flow, an extra term (the second term on the right hand*

side of (2.2) *in Proposition 2.1) appears due to the fact that the metrics are evolving. In order to make the maximum principle argument work, it is required that* $b > 1$. *Moreover, our method used in the proofs of Theorem 2.6 and Theorem 2.11 can also be applied to the FDE* (1.1) *in the fixed metric case to prove certain Aronson-Bénilan estimates with* $b > 1$ *and* $p \in (1 - \frac{2(b-1)}{bn}, 1)$.

Furthermore, we apply similar techniques used in the proof of Theorem 1.1 to extend the Li-Yau-Hamilton estimates (1.8) and (1.9), for the conjugate heat equation and the heat equation under the Ricci flow respectively, to the complete noncompact setting.

Theorem 1.3. *Let* $g(t)$, $t \in [0, T)$, *be a complete solution to the Ricci flow on* M^n *with bounded curvature and nonnegative scalar curvature. If* u *is a positive solution to the conjugate heat equation* (1.7), *then for* $v = -\ln u - \frac{n}{2}\ln(4\pi(T-t))$, *we have*

$$2\Delta v - |\nabla v|^2 + R \leq \frac{2n}{T-t}$$

on $M \times [0, T)$.

Theorem 1.4. *Let* $(M^n, g_{ij}(t))$, $t \in [0, T)$, *be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator. If* u *is a positive solution to the heat equation* (1.10) *and* $v = -\ln u$, *then we have*

$$2\Delta v - |\nabla v|^2 - 3R \leq \frac{2n}{t} \tag{1.12}$$

on $M \times (0, T)$.

The proofs of Theorem 1.3 and Theorem 1.4 are given in Section 3 and Section 4, respectively.

Ackownledgements: Part of the work was carried out while the first author was visiting the University of Macau, where he was partially supported by Science and Technology Development Fund (Macao S.A.R.) Grant FDCT/016/2013/A1, as well as the Projects RDG010 and MYRG2015-00235-FST of University of Macau . The second author would like to thank Professors Qing Ding, Jixiang Fu, Jiaxing Hong, Jun Li, Quanshui Wu, Qi S. Zhang and Weiping Zhang for their support and encouragement. The second author also wants to thank the Shanghai Center for Mathematical Sciences, where part of the work was carried out, for its hospitality.

2. Aronson-Bénilan and Li-Yau-Hamilton estimates for FDE

In this section we study the FDE (1.11) (with $0 < p < 1$) coupled with the Ricci flow (1.5), and prove Aronson-Bénilen and Li-Yau-Hamilton type estimates for its positive solutions.

Suppose $u = u(x, t)$ is a positive solution to the FDE (1.11), with $0 < p < 1$. Notice that the function $v = \frac{p}{p-1}u^{p-1}$ satisfies the equation

$$\frac{\partial v}{\partial t} = (p-1)v\Delta v + |\nabla v|^2 + (p-1)Rv. \tag{2.1}$$

Let

$$F = \frac{|\nabla v|^2}{v} - b\frac{v_t}{v} + (1-b)\frac{R}{v}.$$

If we denote by

$$y = \frac{|\nabla v|^2}{v} + \frac{R}{v} \qquad \text{and} \qquad z = \frac{v_t}{v} + \frac{R}{v},$$

then we have

$$F = y - bz.$$

Next we recall the following result from [7] (see Proposition 2.2 in [7]):

Proposition 2.1. *Suppose u is a smooth positive solution to (1.11) and $v = \frac{p}{p-1}u^{p-1}$. Let*

$$\mathcal{L} = \frac{\partial}{\partial t} - (p-1)v\Delta.$$

Then,

$$\begin{aligned}\mathcal{L}(F) = {} & 2p\nabla_i F \nabla_i v - [\frac{b-1}{v} + p - 1](\frac{\partial R}{\partial t} - 2\nabla_i R \nabla_i v + 2R_{ij}\nabla_i v \nabla_j v) \\ & - 2(p-1)|\nabla^2 v + \frac{b}{2}Rc|^2 + \frac{(b-2)^2}{2}(p-1)|Rc|^2 - \frac{1}{b}F^2 \\ & - [(p-1)R + \frac{2(b-1)}{b}\frac{R}{v}]F - \frac{b-1}{b}y^2 - \frac{(b-1)(b-2)}{b}y\frac{R}{v}.\end{aligned} \tag{2.2}$$

We also need the following trace version of Hamilton's matrix Li-Yau estimate for the Ricci flow.

Theorem 2.2 (**Hamilton** [19])**.** *Let $(M^n, g_{ij}(t))$, $t \in [0,T)$, be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator, then for any 1-form V_i on M^n, we have*

$$\frac{\partial R}{\partial t} + 2\nabla_i R V_i + 2R_{ij}V_iV_j + \frac{R}{t} \geq 0$$

for $t \in (0,T)$.

We remark that by the work of S. Brendle in [4], the above result remains valid if we replace the assumption of nonnegative curvature operator by the weaker condition that $(M^n, g(t)) \times \mathbb{R}^2$ has nonnegative isotropic curvature.

Similar to the arguments in [7], we first establish a local differential Harnack estimate which will lead to a global upper bound for our Harnack quantity. Then, refining the existing Harnack inequality with Hamilton's distance like function gives us the final result.

Since the last term in (2.2) has different signs for $b \geq 2$ and $1 < b < 2$, we treat these two cases separately.

2.1. **The case of $1 < b < 2$.** The local differential Harnack inequality in this case is as follows:

Proposition 2.3. *Let $(M^n, g_{ij}(t))$, $t \in [0,T]$, be a complete solution to the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator. If u is a positive solution to (1.11) with $p \in (1 - \frac{2(b-1)}{bn}, 1)$, then for $v = \frac{p}{p-1}u^{p-1}$, any point $O \in M$, any constants $R_0 > 0$, $\delta > 0$, and $b \in (1,2)$, we have*

$$\frac{|\nabla v|^2}{v} - b\frac{v_t}{v} - (b-1)\frac{R}{v} \geq -\frac{b\alpha}{t}\left[1 + \frac{1}{2}\sqrt{\frac{1+\delta}{\delta}} + C_1\frac{t\bar{v}_{max}}{R_0^2} + C_2(\delta)tR_{max}\right]$$

on $\coprod_{t\in(0,T]} B_t(O,R_0) \times \{t\}$, where $C_1 = (8n+128)(1-p) + \frac{32b\alpha p^2}{b-1}$, $C_2(\delta) = 17 - p + \sqrt{\frac{b\alpha(2-b)^2(1-p)}{2} + \delta(1-p)^2}$, $\alpha = \frac{bn(1-p)}{2-bn(1-p)}$, $\bar{v}_{max} = \sup_{\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}}(-v)$, and $R_{max} = \sup_{M\times[0,T]} R(x,t)$.

Proof. From (2.2), we have

$$\begin{aligned}\mathcal{L}(F) &= 2p\nabla_i F\nabla_i v + [\frac{b-1}{-v} + 1 - p](\frac{\partial R}{\partial t} - 2\nabla_i R\nabla_i v + 2R_{ij}\nabla_i v\nabla_j v)\\ &\quad + 2(1-p)|\nabla^2 v + \frac{b}{2}Rc|^2 - \frac{(b-2)^2}{2}(1-p)|Rc|^2 - \frac{1}{b}(y-bz)^2\\ &\quad + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]F - \frac{b-1}{b}y^2 + \frac{(b-1)(2-b)}{b}\cdot\frac{R}{v}y.\end{aligned}$$

Let $\eta(s)$ be a smooth function defined for $s \geq 0$ such that $\eta(s) = 1$ for $0 \leq s \leq \frac{1}{2}$, $\eta(s) = 0$ for $s \geq 1$, $\eta(s) > 0$ for $\frac{1}{2} < s < 1$, $|\eta'| \leq 16\eta^{\frac{1}{2}}$ and $\eta'' \geq -16\eta \geq -16$. Define

$$\phi(x,t) = \eta\left(\frac{r(x,t)}{2R_0}\right) \tag{2.3}$$

on $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, where $r(x,t)$ is the distance function from O at time t. It follows that

$$\begin{aligned}t\phi\mathcal{L}(t\phi F) &= t\phi^2 F + t^2\phi F\phi_t - (p-1)t^2 v\phi F\Delta\phi - 2(p-1)t^2 v\phi\nabla_i\phi\nabla_i F + t^2\phi^2\mathcal{L}(F)\\ &= \phi(\tilde{y} - b\tilde{z}) + t\phi_t(\tilde{y} - b\tilde{z}) - (p-1)tv\Delta\phi(\tilde{y} - b\tilde{z}) - 2(p-1)t^2 v\phi\nabla_i\phi\nabla_i F\\ &\quad + 2pt^2\phi^2\nabla_i F\nabla_i v + t\phi^2[\frac{b-1}{-v} + 1 - p]Q - t\phi^2 R[\frac{b-1}{-v} + 1 - p]\\ &\quad + 2(1-p)t^2\phi^2|\nabla^2 v + Rc - \frac{2-b}{2}Rc|^2 - \frac{(b-2)^2}{2}(1-p)t^2\phi^2|Rc|^2 - \frac{1}{b}(\tilde{y} - b\tilde{z})^2\\ &\quad + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]t\phi(\tilde{y} - b\tilde{z}) - \frac{b-1}{b}\tilde{y}^2 + \frac{(b-1)(2-b)}{b}\cdot\frac{R}{v}t\phi\tilde{y},\end{aligned}$$

where $\tilde{y} = t\phi y$, $\tilde{z} = t\phi z$, and

$$Q = t\frac{\partial R}{\partial t} + R - 2t\nabla_i R\nabla_i v + 2tR_{ij}\nabla_i v\nabla_j v \geq 0. \tag{2.4}$$

Note that inequality (2.4) is a consequence of Theorem 2.2.

It was shown in [7] that

$$|\frac{\partial\phi}{\partial t}| \leq 16R_{max}, \qquad \Delta\phi \geq -\frac{8n}{R_0^2}, \qquad |\nabla\phi| \leq \frac{8}{R_0}\phi^{\frac{1}{2}}.$$

If $t\phi F \geq 0$ in $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, then the theorem is automatically true. Otherwise, since $t\phi F = 0$ on the parabolic boundary of $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, we may assume that $t\phi F$ achieves a negative minimum at time $t_0 > 0$ and some interior point x_0. Thus, at (x_0,t_0), we have

$$\tilde{y} - b\tilde{z} = t_0\phi F < 0, \qquad F\nabla\phi = -\phi\nabla F, \qquad \text{and} \qquad \mathcal{L}(t\phi F)(x_0,t_0) \leq 0.$$

Moreover, since

$$-2pt_0^2\phi F\nabla_i\phi\nabla_i v \geq 2pt_0^2\phi F|\nabla\phi||\nabla v| \geq t_0\phi F\frac{16p}{R_0}(-\tilde{y})^{\frac{1}{2}}(-t_0 v)^{\frac{1}{2}},$$

$$R + \Delta v = y - z, \qquad \text{and} \qquad -\tilde{y} \geq t\phi\frac{R}{-v},$$

by the Cauchy-Schwarz inequality, we have

$$
\begin{aligned}
0 \geq & t_0\phi\mathcal{L}(t\phi F) \\
\geq & (\tilde{y}-b\tilde{z})+16t_0R_{max}(\tilde{y}-b\tilde{z})+(8n+128)\frac{(p-1)t_0v}{R_0^2}(\tilde{y}-b\tilde{z})+\frac{16p}{R_0}(-\tilde{y})^{\frac{1}{2}}(-t_0v)^{\frac{1}{2}}(\tilde{y}-b\tilde{z}) \\
& -(1-p)t_0\phi R-(b-1)t_0\phi(\frac{R}{-v})+\frac{2}{b^2n(1-p)}[\tilde{y}-b\tilde{z}-(b-1)(-\tilde{y})-\frac{b(2-b)}{2}(1-p)t_0\phi R]^2 \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2\phi^2R^2-\frac{1}{b}(\tilde{y}-b\tilde{z})^2+[(1-p)R+\frac{2(b-1)}{b}\frac{R}{-v}]t\phi(\tilde{y}-b\tilde{z}) \\
& -\frac{b-1}{b}\tilde{y}^2+\frac{(b-1)(2-b)}{b}\cdot t_0^2\phi^2(\frac{R}{-v})^2 \\
\geq & (\tilde{y}-b\tilde{z})+16t_0R_{max}(\tilde{y}-b\tilde{z})+(8n+128)\frac{(p-1)t_0v}{R_0^2}(\tilde{y}-b\tilde{z})+\frac{16p}{R_0}(-\tilde{y})^{\frac{1}{2}}(-t_0v)^{\frac{1}{2}}(\tilde{y}-b\tilde{z}) \\
& -(1-p)t_0\phi R+\frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2+\frac{b-1}{b}[\frac{2(b-1)}{bn(1-p)}-1](-\tilde{y})^2-\frac{4(b-1)}{b^2n(1-p)}(-\tilde{y})(\tilde{y}-b\tilde{z}) \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2\phi^2R^2+[(1-p)R+\frac{2(b-1)}{b}\frac{R}{-v}]t_0\phi(\tilde{y}-b\tilde{z}) \\
& +\frac{(b-1)(2-b)}{b}\cdot t^2\phi^2(\frac{R}{-v})^2-(b-1)t\phi(\frac{R}{-v}).
\end{aligned}
$$

According to the assumption on p, we have $\frac{2(b-1)}{bn(1-p)}-1\geq 0$. Moreover, if we choose $\beta=\frac{2(b-1)}{b\alpha}$, then we have

$$
\begin{aligned}
0 \geq & (\tilde{y}-b\tilde{z})+(16+(1-p))t_0R_{max}(\tilde{y}-b\tilde{z})+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}(\tilde{y}-b\tilde{z}) \\
& -(1-p)t_0\phi R+\frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2-(\frac{2(b-1)}{b\alpha}-\beta)(-\tilde{y})(\tilde{y}-b\tilde{z}) \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2\phi^2R^2+\frac{(b-1)^2}{b\alpha}\cdot t_0^2\phi^2(\frac{R}{-v})^2-(b-1)t_0\phi(\frac{R}{-v}) \\
\geq & \frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2+\left[1+(17-p)t_0R_{max}+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}\right](\tilde{y}-b\tilde{z}) \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2R_{max}^2-(1-p)t_0R_{max}-\frac{b\alpha}{4}.
\end{aligned}
$$

This implies that

$$
\begin{aligned}
\tilde{y}-b\tilde{z} \geq & -b\alpha\left[1+(17-p)t_0R_{max}+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}\right] \\
& -\sqrt{b\alpha\left[\frac{(b-2)^2}{2}(1-p)t_0^2R_{max}^2+(1-p)t_0R_{max}+\frac{b\alpha}{4}\right]}.
\end{aligned}
$$

Now Proposition 2.3 follows easily. □

If u is bounded, then letting $R_0\to\infty$, we get a global differential Harnack inequality.

Corollary 2.4. *Let $(M^n, g_{ij}(t))$, $t\in[0,T]$, be a complete solution to the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator. If u is a smooth positive bounded*

solution to (1.2) *with* $1-\frac{2(b-1)}{bn}<p<1$, *then for* $v=\frac{p}{p-1}u^{p-1}$, *any constants* $\delta>0$, *and* $b\in(1,2)$, *we have*

$$\frac{|\nabla v|^2}{v}-b\frac{v_t}{v}-(b-1)\frac{R}{v}\geq-\frac{b\alpha}{t}\left[1+\frac{1}{2}\sqrt{\frac{1+\delta}{\delta}}+C_2(\delta)tR_{max}\right]$$

on $M\times(0,T]$, *where* $C_2(\delta)$, α, *and* R_{max} *are the same constants as in Proposition 2.3.*

We shall need the following lemma about some distance-like function constructed by Hamilton (see e.g. [19] and [8]),

Lemma 2.5. *Let* $g_{ij}(t)$, $t\in[0,T]$, *be a complete solution to the Ricci flow on* M^n *with bounded curvature tensor. Then, there exists a smooth function* $f(x)$ *on* M *and a positive constant* $C>0$ *such that* $f\geq 1$, $f(x)\to\infty$ *as* $d_0(x,O)\to\infty$ *(for any fixed point* $O\in M$*),*

$$|\nabla f|_{g(t)}\leq C,\qquad and\qquad |\nabla\nabla f|_{g(t)}\leq C$$

on $M\times[0,T]$.

Now using the function f in the above Lemma and a method of Hamilton, we are able to refine the differential Harnack inequality in Corollary 2.4.

Theorem 2.6. *Let* $(M^n,g_{ij}(t))$, $t\in[0,T]$, *be a complete solution to the Ricci flow* (1.5) *with bounded curvature and nonnegative curvature operator. If* u *is a smooth positive bounded solution to* (1.11) *with* $1-\frac{2(b-1)}{bn}<p<1$, *then for* $v=\frac{p}{p-1}u^{p-1}$, *and* $b\in(1,2)$, *we have*

$$\frac{|\nabla v|^2}{v}-b\frac{v_t}{v}-(b-1)\frac{R}{v}+KR_{max}+\frac{b\alpha}{t}\geq 0$$

on $M\times(0,T]$, *where* $\alpha=\frac{bn(1-p)}{2-bn(1-p)}$, $K=\max\{\frac{(b\alpha+1)(1-p)}{2},b\alpha(1-p)+(2-b)\sqrt{\frac{b\alpha(n-1)(1-p)}{2n}}\}$, *and* $R_{max}=\sup_{M\times[0,T]}R(x,t)$.

Proof. From (2.2), we have

$$\begin{aligned}\mathcal{L}(F)=&\,2p\nabla_iF\nabla_iv+[\frac{b-1}{-v}+1-p](\frac{\partial R}{\partial t}-2\nabla_iR\nabla_iv+2R_{ij}\nabla_iv\nabla_jv)\\&+2(1-p)|\nabla^2v+\frac{b}{2}Rc|^2-\frac{(b-2)^2}{2}(1-p)|Rc|^2-\frac{1}{b}F^2\\&+[(1-p)R+\frac{2(b-1)}{b}\frac{R}{-v}]F-\frac{b-1}{b}y^2+\frac{(b-1)(2-b)}{b}\cdot\frac{R}{v}y.\end{aligned}$$

Let $H = t(F+K)+d$, then

$$\begin{aligned}
t\mathcal{L}(H) &= t(F+K) + t^2\mathcal{L}(F)\\
&= H - d + 2pt\nabla_i H\nabla_i v + t[\frac{b-1}{-v}+1-p]Q - tR[\frac{b-1}{-v}+1-p]\\
&\quad + 2(1-p)t^2|\nabla^2 v + Rc - \frac{2-b}{2}Rc|^2 - \frac{(b-2)^2}{2}(1-p)t^2|Rc|^2 - \frac{1}{b}(H-tK-d)^2\\
&\quad + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]t(H-tK-d) - \frac{b-1}{b}(ty)^2 + \frac{(b-1)(2-b)}{b}\cdot\frac{t^2R}{v}y\\
&\geq H - d + 2pt\nabla_i H\nabla_i v - tR[\frac{b-1}{-v}+1-p]\\
&\quad + \frac{2}{b^2n(1-p)}\left[H - tK - d - (b-1)(-ty) - \frac{b(2-b)}{2}(1-p)tR\right]^2\\
&\quad - \frac{(b-2)^2}{2}(1-p)t^2R^2 - \frac{1}{b}(H-tK-d)^2 + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]t(H-tK-d)\\
&\quad - \frac{b-1}{b}(ty)^2 + \frac{(b-1)(2-b)}{b}\cdot(\frac{tR}{-v})(-ty)\\
&\geq H - d + 2pt\nabla_i H\nabla_i v + (b-1)(-\frac{2d}{b}-1)\frac{tR}{-v} + [\frac{2(2-b)d}{bn(1-p)} - d - 1](1-p)tR\\
&\quad + \frac{1}{b\alpha}(H-tK-d)^2 + \frac{b-1}{b}[\frac{2(b-1)}{bn(1-p)}-1](-ty)^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t^2R^2\\
&\quad - \frac{4(b-1)}{b^2n(1-p)}(-ty)(H-tK-d) + \frac{2(b-1)}{b}\frac{tR}{-v}(H-tK)\\
&\quad - [\frac{2(2-b)}{bn(1-p)}-1](1-p)tR(H-tK).
\end{aligned}$$

Let $\tilde{H} = H + \epsilon\psi$, where $\psi = e^{At}f$ for some constant $A > 0$ to be determined, and f being the function in Lemma 2.5. Then

$$t\mathcal{L}(\tilde{H}) = t\mathcal{L}(H) + tA\epsilon\psi - (p-1)vt\epsilon e^{At}\Delta f.$$

From Corollary 2.4, we know that $\tilde{H} > 0$ at $t = 0$ and outside a fixed compact subset of M for $t \in (0, T]$. Suppose that $\tilde{H}$ reaches 0 for the first time at some point $x_0 \in M$ when $t = t_0 > 0$. Then we have at (x_0, t_0),

$$H = -\epsilon\psi < 0, \qquad \nabla\tilde{H} = 0, \qquad \text{and} \qquad \mathcal{L}(\tilde{H}) \leq 0.$$

Setting $d = b\alpha$, and $\beta = \frac{2(b-1)}{bn(1-p)} - 1 > 0$ we have

$$\begin{aligned}
0 \geq& t_0\mathcal{L}(\tilde{H})\\
\geq& -\epsilon\psi - b\alpha - 2pt_0\epsilon e^{At}\nabla_i f\nabla_i v + [\frac{2(2-b)b\alpha}{bn(1-p)} - b\alpha - 1](1-p)t_0R + \frac{1}{b\alpha}(\epsilon\psi + t_0K + b\alpha)^2\\
&+ \frac{\beta(b-1)}{b}(-t_0y)^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t_0^2R^2\\
&+ [\frac{2(2-b)}{bn(1-p)} - 1](1-p)t_0R(\epsilon\psi + t_0K) + t_0A\epsilon\psi - (p-1)vt_0\epsilon e^{At_0}\Delta f
\end{aligned}$$

$$
\begin{aligned}
&\geq -Ct_0\epsilon\psi|\nabla v| + [\frac{2(2-b)b\alpha}{bn(1-p)} - b\alpha - 1](1-p)t_0R + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{1}{b\alpha}t_0^2K^2 + \epsilon\psi + \frac{2}{b\alpha}t_0K\epsilon\psi \\
&\quad + 2t_0K + \frac{\beta(b-1)}{b}(\frac{t_0|\nabla v|^2}{-v})^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t_0^2R^2 \\
&\quad + [\frac{2(2-b)}{bn(1-p)} - 1](1-p)t_0R(\epsilon\psi + t_0K) + t_0A\epsilon\psi - C\bar{v}_{max}t_0\epsilon\psi \\
&\geq -\delta\epsilon\psi\frac{t_0|\nabla v|^2}{-v} - \frac{C}{4\delta}t_0(-v)\epsilon\psi + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{\beta(b-1)}{b}(\frac{t_0|\nabla v|}{-v})^2 \\
&\quad - [b\alpha+1](1-p)t_0R + 2t_0K + \frac{1}{b\alpha}t_0^2K^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t_0^2R^2 \\
&\quad - (1-p)t_0^2RK + \left(A + \frac{2}{b\alpha}K + [\frac{2(2-b)}{bn(1-p)} - 1](1-p)R - C\bar{v}_{max}\right)t_0\epsilon\psi.
\end{aligned}
$$

We may choose a suitable δ so that

$$-\delta\epsilon\psi\frac{t_0|\nabla v|^2}{-v} + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{\beta(b-1)}{b}(\frac{t_0|\nabla v|^2}{-v})^2 \geq 0.$$

Notice that by choosing $K = \max\{\frac{(b\alpha+1)(1-p)}{2}, b\alpha(1-p) + (2-b)\sqrt{\frac{b\alpha(n-1)(1-p)}{2n}}\}R_{max}$, we can make both

$$-[b\alpha+1](1-p)R + 2K \geq 0,$$

and

$$\frac{1}{b\alpha}K^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)R^2 - (1-p)RK \geq 0.$$

Hence if we choose $A > (\frac{C}{4\delta} + C)\bar{v}_{max} + (1-p)R_{max}$, then we have

$$
\begin{aligned}
0 \geq& t\mathcal{L}(\tilde{H}) \\
\geq& \left[A + \frac{2}{b\alpha}K - (1-p)R - (C + \frac{C}{4\delta})\bar{v}_{max}\right]t_0\epsilon\psi \\
>& 0,
\end{aligned}
$$

a contradiction. Therefore, $\tilde{H} > 0$ for all $t \in [0,T]$. Letting $\epsilon \to 0$, we get

$$H \geq 0.$$

□

As in [7], consequently we have the following Harnack inequalities.

Corollary 2.7. *Suppose that $(M, g_{ij}(t))$, $t \in [0,T]$, is a complete solution of the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator, and u is a bounded smooth positive solution to (1.11) for $1 - \frac{2(b-1)}{bn} < p < 1$ and $b \in (1,2)$. Let $\bar{v} = -v = \frac{p}{1-p}u^{p-1}$, if $\bar{v}_{min} = \inf_{M\times[0,T]}\bar{v} > 0$, then for any points x_1,$x_2 \in M$ and $0 < t_1 < t_2$, we have*

$$\bar{v}(x_2,t_2) \leq \bar{v}(x_1,t_1)\cdot(\frac{t_2}{t_1})^\alpha exp\left(\frac{\Gamma}{\bar{v}_{min}} + \frac{KR_{max}}{b}(t_2-t_1)\right),$$

where α, K and R_{max} are the same constants as in Theorem 2.6, and $\Gamma = \inf_\gamma\int_{t_1}^{t_2}(\frac{b-1}{b}R + \frac{b}{4}\left|\frac{d\gamma}{d\tau}\right|^2_{g_{ij}(\tau)})d\tau$ with the infimum taking over all smooth curves $\gamma(\tau)$ in M, $\tau \in [t_1,t_2]$, with $\gamma(t_1) = x_1$ and $\gamma(t_2) = x_2$.

Corollary 2.8. *Suppose that $(M, g_{ij}(t))$, $t \in [0,T]$, is a complete solution of the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator, and u is a bounded smooth positive solution to (1.11) for $1 - \frac{2(b-1)}{bn} < p < 1$ and $b \in (1,2)$. Let $\bar{v} = -v = \frac{p}{1-p}u^{p-1}$, then for any points x_1,$x_2 \in M$ and $0 < t_1 < t_2$, we have*

$$\bar{v}(x_2,t_2) - \bar{v}(x_1,t_1) \le \alpha \bar{v}_{max} \ln \frac{t_2}{t_1} + [\frac{b-1}{b} + \frac{K}{b}\bar{v}_{max}]R_{max}(t_2 - t_1) + \frac{b}{4}\frac{d^2_{t_1}(x_1,x_2)}{t_2 - t_1}$$

where α, K and R_{max} are the same constants as in Theorem 2.6, and $\bar{v}_{max} = \sup_{M\times[0,T]} \bar{v}$.

2.2. **The case of $b \ge 2$.** In this case, we first have

Proposition 2.9. *Let $(M^n, g_{ij}(t))$, $t \in [0,T]$, be a complete solution to the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator. If u is a smooth positive solution to (1.11) with $p \in (1-\frac{2}{bn}, 1)$ and $b \ge 2$, then for $v = \frac{p}{p-1}u^{p-1}$, any point $O \in M$, any constants $R_0 > 0$, $\delta_1 \in (0,1)$ and $\delta_2 > 0$, we have*

$$\frac{|\nabla v|^2}{v} - b\frac{v_t}{v} - (b-1)\frac{R}{v} \ge -\frac{b\alpha}{t}\left[1 + \frac{1}{2}\sqrt{\frac{1}{1-\delta_1} + \delta_2} + C_1(\delta_1,\delta_2)R_{max} + C_2\frac{t\bar{v}_{max}}{R_0^2}\right]$$

on $\coprod_{t\in(0,T]} B_t(O,R_0)\times\{t\}$, where

$$C_1(\delta_1,\delta_2) = 17 - p + \frac{2(b-2)}{bn} + \sqrt{b\alpha\left[\frac{(2-b)^2(1-p)}{2} + \frac{\alpha(b-2)^2}{bn^2\delta_1}\right] + \frac{(1-p)^2}{\delta_2}},$$

$$C_2 = (8n+128)(1-p) + \frac{32b\alpha p^2}{b-1},$$

$\alpha = \frac{bn(1-p)}{2-bn(1-p)}$, $\bar{v}_{max} = \max_{\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}}\{-v\}$ *and* $R_{max} = \sup_{M\times[0,T]} R$.

Proof. Let $\phi(x,t)$ be the same cut-off function as in (2.3). It follows from (2.2) that

$$\begin{aligned} t\phi\mathcal{L}(t\phi F) &= \phi(\tilde{y} - b\tilde{z}) + t\phi_t(\tilde{y} - b\tilde{z}) - (p-1)tv\Delta\phi(\tilde{y} - b\tilde{z}) - 2(p-1)t^2v\phi\nabla_i\phi\nabla_i F \\ &\quad + 2pt^2\phi^2\nabla_i F\nabla_i v + t\phi^2[\frac{b-1}{-v} + 1 - p]Q - t\phi^2 R[\frac{b-1}{-v} + 1 - p] \\ &\quad + 2(1-p)t^2\phi^2|\nabla^2 v + Rc + \frac{b-2}{2}Rc|^2 - \frac{(b-2)^2}{2}(1-p)t^2\phi^2|Rc|^2 - \frac{1}{b}(\tilde{y} - b\tilde{z})^2 \\ &\quad + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]t\phi(\tilde{y} - b\tilde{z}) - \frac{b-1}{b}\tilde{y}^2 - \frac{(b-1)(b-2)}{b}\cdot\frac{R}{v}t\phi\tilde{y}, \end{aligned}$$

where $\tilde{y} = t\phi y$, $\tilde{z} = t\phi z$, and $Q \ge 0$ is the quantity in (2.4).

If $t\phi F \ge 0$ in $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, then we are done. Otherwise, since $t\phi F = 0$ on the parabolic boundary of $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, we may assume that $t\phi F$ achieves a negative minimum for the first time at $t_0 > 0$ and some interior point x_0. Thus, at (x_0,t_0), we have

$$\tilde{y} - b\tilde{z} = t_0\phi F < 0, \qquad \nabla\phi = -\phi\nabla F, \qquad \text{and} \qquad \mathcal{L}(t\phi F)(x_0,t_0) \le 0.$$

Moreover, since

$$-2pt_0^2\phi F\nabla_i\phi\nabla_i v \ge 2pt_0^2\phi F|\nabla\phi||\nabla v| \ge t_0\phi F\frac{16p}{R_0}(-\tilde{y})^{\frac{1}{2}}(-t_0 v)^{\frac{1}{2}},$$

we get

$$
\begin{aligned}
0 \geq & t_0\phi\mathcal{L}(t\phi F) \\
\geq & (\tilde{y}-b\tilde{z})+16t_0R_{max}(\tilde{y}-b\tilde{z})+(8n+128)\frac{(p-1)t_0v}{R_0^2}(\tilde{y}-b\tilde{z})+\frac{16p}{R_0}(-\tilde{y})^{\frac{1}{2}}(-t_0v)^{\frac{1}{2}}(\tilde{y}-b\tilde{z}) \\
& -(1-p)t_0\phi R-(b-1)t_0\phi(\frac{R}{-v})+\frac{2}{b^2n(1-p)}[\tilde{y}-b\tilde{z}-(b-1)(-\tilde{y})+\frac{b(b-2)}{2}(1-p)t_0\phi R]^2 \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2\phi^2R^2-\frac{1}{b}(\tilde{y}-b\tilde{z})^2+[(1-p)R+\frac{2(b-1)}{b}\frac{R}{-v}]t_0\phi(\tilde{y}-b\tilde{z}) \\
& -\frac{b-1}{b}\tilde{y}^2-\frac{(b-1)(b-2)}{b}\cdot(-\tilde{y})^2 \\
\geq & (\tilde{y}-b\tilde{z})+16t_0R_{max}(\tilde{y}-b\tilde{z})+(8n+128)\frac{(p-1)t_0v}{R_0^2}(\tilde{y}-b\tilde{z})+\beta(-\tilde{y})(\tilde{y}-b\tilde{z}) \\
& +\frac{64p^2}{\beta R_0^2}(-t_0v)(\tilde{y}-b\tilde{z})-(1-p)t_0\phi R+\frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2+\frac{(b-1)^2}{b\alpha}(-\tilde{y})^2-(b-1)t\phi(\frac{R}{-v}) \\
& -\frac{2(b-1)(b-2)}{bn}t_0\phi R(-\tilde{y})-\frac{4(b-1)}{b^2n(1-p)}(-\tilde{y})(\tilde{y}-b\tilde{z})+\frac{2(b-2)}{bn(1-p)}(1-p)t_0\phi R(\tilde{y}-b\tilde{z}) \\
& -\frac{(b-2)^2}{2}(1-p)t_0^2\phi^2R^2+[(1-p)R+\frac{2(b-1)}{b}\frac{R}{-v}]t_0\phi(\tilde{y}-b\tilde{z}).
\end{aligned}
$$

If we choose $\beta=\frac{2(b-1)}{b\alpha}$ and $0<\delta_1<1$, then we have

$$
\begin{aligned}
0 \geq & (\tilde{y}-b\tilde{z})+(17-p+\frac{2(b-2)}{bn})t_0R_{max}(\tilde{y}-b\tilde{z})+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}(\tilde{y}-b\tilde{z}) \\
& -(1-p)t_0\phi R+\frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2-(\frac{2(b-1)}{b\alpha}-\beta)(-\tilde{y})(\tilde{y}-b\tilde{z})+\frac{(b-1)^2}{b\alpha}(1-\delta_1)\cdot t^2\phi^2(\frac{R}{-v})^2 \\
& -(b-1)t\phi(\frac{R}{-v})-\left[\frac{(b-2)^2}{2}(1-p)+\frac{\alpha(b-2)^2}{bn^2\delta_1}\right]t_0^2\phi^2R^2 \\
\geq & \frac{1}{b\alpha}(\tilde{y}-b\tilde{z})^2+\left[1+(17-p+\frac{2(b-2)}{bn})t_0R_{max}+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}\right](\tilde{y}-b\tilde{z}) \\
& -\left[\frac{(b-2)^2}{2}(1-p)+\frac{\alpha(b-2)^2}{bn^2\delta_1}\right]t_0^2R_{max}^2-(1-p)t_0R_{max}-\frac{b\alpha}{4(1-\delta_1)}.
\end{aligned}
$$

Therefore, we have

$$
\begin{aligned}
\tilde{y}-b\tilde{z} \geq & -b\alpha\left[1+(17-p+\frac{2(b-2)}{bn})t_0R_{max}+[(8n+128)(1-p)+\frac{64p^2}{\beta}]\frac{t_0\bar{v}_{max}}{R_0^2}\right] \\
& -\sqrt{b\alpha\left(\left[\frac{(b-2)^2}{2}(1-p)+\frac{\alpha(b-2)^2}{bn^2\delta_1}\right]t_0^2R_{max}^2+(1-p)t_0R_{max}+\frac{b\alpha}{4(1-\delta_1)}\right)}.
\end{aligned}
$$

This finishes the proof. □

Assuming further that u is bounded and letting $R_0\to\infty$, we have

Corollary 2.10. *Let $(M^n, g_{ij}(t))$, $t\in[0,T]$, be a complete solution to the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator. If u is a bounded smooth positive*

solution to (1.11) *with* $p \in (1-\frac{2}{bn}, 1)$ *and* $b \geq 2$, *then for* $v = \frac{p}{p-1}u^{p-1}$, *and any constants* $\delta_1 \in (0,1)$ *and* $\delta_2 > 0$, *we have*

$$\frac{|\nabla v|^2}{v} - b\frac{v_t}{v} - (b-1)\frac{R}{v} \geq -\frac{b\alpha}{t}\left[1 + \frac{1}{2}\sqrt{\frac{1}{1-\delta_1} + \delta_2} + C_1(\delta_1, \delta_2)R_{max}\right]$$

on $M \times (0,T]$, *where* α, $C_1(\delta_1, \delta_2)$ *and* R_{max} *are the same constants as in Proposition 2.9.*

Using Hamilton's method, again we can obtain the following refined estimate.

Theorem 2.11. *Let* $(M^n, g_{ij}(t))$, $t \in [0,T]$, *be a complete solution to the Ricci flow* (1.5) *with bounded curvature and nonnegative curvature operator. If* u *is a bounded smooth positive solution to* (1.11) *with* $p \in (1-\frac{2}{bn}, 1)$ *and* $b \geq 2$, *then for* $v = \frac{p}{p-1}u^{p-1}$, *we have*

$$\frac{|\nabla v|^2}{v} - b\frac{v_t}{v} - (b-1)\frac{R}{v} + KR_{max} + \frac{b\alpha}{t} \geq 0$$

on $M \times (0,T]$, *where* α *and* R_{max} *are the same constants as in Proposition 2.9, and*

$$K = \max\{[\frac{2(b-2)\alpha}{n(1-p)} + b\alpha + 1]\frac{(1-p)}{2},\ b\alpha[\frac{2(b-2)}{bn(1-p)} + 1](1-p) + (b-2)\sqrt{\frac{b\alpha(n-1)(1-p)}{2n}}\}.$$

Proof. Let $H = t(F+K) + d$ with $d = b\alpha$ and $K > 0$ to be determined. From (2.2), one has

$$\begin{aligned}
t\mathcal{L}(H) \geq & H - d + 2pt\nabla_i H \nabla_i v - tR[\frac{b-1}{-v} + 1 - p] - \frac{b-1}{b}(ty)^2 - \frac{(b-1)(b-2)}{b}\cdot(ty)^2 \\
& + \frac{2}{b^2 n(1-p)}\left[H - tK - d - (b-1)(-ty) + \frac{b(b-2)}{2}(1-p)tR\right]^2 \\
& - \frac{(b-2)^2}{2}(1-p)t^2R^2 - \frac{1}{b}(H - tK - d)^2 + [(1-p)R + \frac{2(b-1)}{b}\frac{R}{-v}]t(H - tK - d) \\
\geq & H - d + 2pt\nabla_i H \nabla_i v + (b-1)(-\frac{2d}{b} - 1)\frac{tR}{-v} - [\frac{2(b-2)d}{bn(1-p)} + d + 1](1-p)tR \\
& + \frac{1}{b\alpha}(H - tK - d)^2 + \frac{(b-1)^2}{b\alpha}(-ty)^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t^2R^2 \\
& - \frac{4(b-1)}{b^2 n(1-p)}(-ty)(H - tK - d) + \frac{2(b-1)}{b}\frac{tR}{-v}(H - tK) \\
& + [\frac{2(b-2)}{bn(1-p)} + 1](1-p)tR(H - tK) - \frac{2(b-1)(b-2)}{bn}tR(-ty).
\end{aligned}$$

Let $\tilde{H} = H + \epsilon\psi$, where $\psi = e^{At}f$ for some constant $A > 0$ to be determined, and f being the function in Lemma 2.5. From Corollary 2.10, we know that $\tilde{H} > 0$ at $t = 0$ and outside a fixed compact subset of M for $t \in (0,T]$. Suppose that $\tilde{H}$ reaches 0 for the first time at some point $x_0 \in M$ when $t = t_0 > 0$. Then we have at (x_0, t_0),

$$H = -\epsilon\psi < 0, \qquad \nabla\tilde{H} = 0, \qquad \text{and} \qquad \mathcal{L}(\tilde{H}) \leq 0.$$

Hence, we get

$$\begin{aligned}
0 \geq & t_0\mathcal{L}(\tilde{H}) \\
\geq & -Ct_0\epsilon\psi|\nabla v| - [\frac{2(b-2)b\alpha}{bn(1-p)} + b\alpha + 1](1-p)t_0R + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{1}{b\alpha}t_0^2K^2 + \epsilon\psi \\
& + \frac{2}{b\alpha}t_0K\epsilon\psi + 2t_0K + \frac{(b-1)^2}{b\alpha}(\frac{t_0|\nabla v|^2}{-v})^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t_0^2R^2 \\
& - [\frac{2(b-2)}{bn(1-p)} + 1](1-p)t_0R(\epsilon\psi + t_0K) + t_0A\epsilon\psi - C\bar{v}_{max}t_0\epsilon\psi \\
& + [\frac{4(b-1)}{b^2n(1-p)}K - \frac{2(b-1)(b-2)}{bn}R](-t_0^2y) \\
\geq & -\delta\epsilon\psi\frac{t_0|\nabla v|^2}{-v} - \frac{C}{4\delta}t_0(-v)\epsilon\psi + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{(b-1)^2}{b\alpha}(\frac{t_0|\nabla v|^2}{-v})^2 \\
& - [\frac{2(b-2)\alpha}{n(1-p)} + b\alpha + 1](1-p)t_0R + 2t_0K + \frac{1}{b\alpha}t_0^2K^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)t_0^2R^2 \\
& - [\frac{2(b-2)}{bn(1-p)} + 1](1-p)t_0^2RK + \left(A + \frac{2}{b\alpha}K - [\frac{2(2-b)}{bn(1-p)} + 1](1-p)R - C\bar{v}_{max}\right)t_0\epsilon\psi \\
& + [\frac{4(b-1)}{b^2n(1-p)}K - \frac{2(b-1)(b-2)}{bn}R](-t_0^2y).
\end{aligned}$$

We may choose appropriate δ so that

$$-\delta\epsilon\psi\frac{t_0|\nabla v|^2}{-v} + \frac{1}{b\alpha}(\epsilon\psi)^2 + \frac{(b-1)^2}{b\alpha}(\frac{t_0|\nabla v|^2}{-v})^2 \geq 0.$$

Notice that by choosing

$$K = \max\{[\frac{2(b-2)\alpha}{n(1-p)} + b\alpha + 1]\frac{(1-p)}{2}, b\alpha[\frac{2(b-2)}{bn(1-p)} + 1](1-p) + (b-2)\sqrt{\frac{b\alpha(n-1)(1-p)}{2n}}\}R_{max},$$

we can make

$$\frac{4(b-1)}{b^2n(1-p)}K - \frac{2(b-1)(b-2)}{bn}R \geq 0,$$

$$-[\frac{2(b-2)\alpha}{n(1-p)} + b\alpha + 1](1-p)R + 2K \geq 0,$$

and

$$\frac{1}{b\alpha}K^2 - \frac{(b-2)^2(n-1)}{2n}(1-p)R^2 - [\frac{2(b-2)}{bn(1-p)} + 1](1-p)RK \geq 0.$$

Hence if we furthermore choose $A > (\frac{C}{4\delta} + C)\bar{v}_{max} + (1-p)R_{max}$, then we get

$$0 \geq t_0\mathcal{L}(\tilde{H}) \geq \left[A - (1-p)R - (C + \frac{C}{4\delta})\bar{v}_{max}\right]t_0\epsilon\psi > 0,$$

a contradiction. Therefore, $\tilde{H} > 0$ for all $t \in [0, T]$. Letting $\epsilon \to 0$, it follows that

$$H \geq 0.$$

□

Integrating the differential Harnack inequality in the above theorem along space-time paths, we obtain

Corollary 2.12. *Suppose that $(M, g_{ij}(t))$, $t \in [0,T]$, is a complete solution of the Ricci flow with bounded curvature and nonnegative curvature operator, and u is a bounded smooth positive solution to (1.11) for $1 - \frac{2}{bn} < p < 1$ and $b \geq 2$. Let $\bar{v} = -v = \frac{p}{1-p}u^{p-1}$, if $\bar{v}_{min} = \inf_{M\times[0,T]} \bar{v} > 0$, then for any points x_1,$x_2 \in M$ and $0 < t_1 < t_2$, we have*

$$\bar{v}(x_2,t_2) \leq \bar{v}(x_1,t_1) \cdot (\frac{t_2}{t_1})^{\alpha} exp\left(\frac{\Gamma}{\bar{v}_{min}} + \frac{KR_{max}}{b}(t_2 - t_1)\right), \tag{2.5}$$

where α, K and R_{max} are the same constants as in Theorem 2.11, and $\Gamma = \inf_{\gamma} \int_{t_1}^{t_2} (\frac{b-1}{b}R + \frac{b}{4}\left|\frac{d\gamma}{d\tau}\right|^2_{g_{ij}(\tau)})d\tau$ with the infimum taking over all smooth curves $\gamma(\tau)$ in M, $\tau \in [t_1,t_2]$, with $\gamma(t_1) = x_1$ and $\gamma(t_2) = x_2$.

Corollary 2.13. *Suppose that $(M, g_{ij}(t))$, $t \in [0,T]$, is a complete solution of the Ricci flow (1.5) with bounded curvature and nonnegative curvature operator, and u is a bounded smooth positive solution to (1.11) for $1 - \frac{2}{bn} < p < 1$ and $b \geq 2$. Let $\bar{v} = -v = \frac{p}{1-p}u^{p-1}$, then for any points x_1,$x_2 \in M$ and $0 < t_1 < t_2$, we have*

$$\bar{v}(x_2,t_2) - \bar{v}(x_1,t_1) \leq \alpha \bar{v}_{max} \ln \frac{t_2}{t_1} + [\frac{b-1}{b} + \frac{K}{b}\bar{v}_{max}]R_{max}(t_2 - t_1) + \frac{b}{4}\frac{d^2_{t_1}(x_1,x_2)}{t_2 - t_1} \tag{2.6}$$

where α, K and R_{max} are the same constants as in Theorem 2.11, and $\bar{v}_{max} = \sup_{M\times[0,T]} \bar{v}$.

3. Li-Yau-Hamilton Estimate for the conjugate Heat equation

In this section, we consider the conjugate heat equation

$$\frac{\partial u}{\partial t} = -\Delta u + Ru \tag{3.1}$$

under the Ricci flow (1.5) and derive Li-Yau type estimates for positive solutions to (3.1). When the underlying manifold is compact, this was first studied indepedently by X. Cao [9] and Kuang-Zhang [23].

Suppose that u is a positive solution of (3.1). Let $u = (4\pi\tau)^{-\frac{n}{2}}e^{-v}$, $\tau = T - t$, Then we have

$$v = -\ln u - \frac{n}{2}\ln(4\pi\tau),$$

and the following evolution equations:

$$\frac{\partial g_{ij}}{\partial \tau} = 2R_{ij},$$

$$\frac{\partial u}{\partial \tau} = \Delta u - Ru,$$

and

$$\frac{\partial v}{\partial \tau} = \Delta v - |\nabla v|^2 + R - \frac{n}{2\tau}.$$

Define

$$\begin{aligned} F &= |\nabla v|^2 - bv_\tau + cR + d\frac{v}{\tau} \\ &= |\nabla v|^2 - b\Delta v + b|\nabla v|^2 - bR + \frac{bn}{2\tau} + cR + d\frac{v}{\tau} \\ &= (b+1)|\nabla v|^2 - b\Delta v + (c-b)R + d\frac{v}{\tau} + \frac{bn}{2\tau}. \end{aligned}$$

Lemma 3.1. *For the operator $\mathcal{L} = \frac{\partial}{\partial\tau} - \Delta$, one has*

$$\mathcal{L}F = -2\nabla_i F\nabla_i v - (2c-b)\Delta R + 2(c+1)\nabla_i R\nabla_i v + \frac{d}{\tau}|\nabla f|^2 - 2(b+2)R_{ij}\nabla_i v\nabla_j v$$
$$- 2|\nabla^2 v|^2 + 2bR_{ij}\nabla_i\nabla_j v - 2(c-b)|Rc|^2 + \frac{d}{\tau}R - \frac{d}{\tau^2}v - \frac{(b+d)n}{2\tau^2}. \tag{3.2}$$

Proof. It is straightforward to check that

$$\begin{aligned}\frac{\partial|\nabla v|^2}{\partial\tau} &= -2R_{ij}\nabla_i v\nabla_j v + 2\nabla_i v_\tau\nabla_i v \\ &= -2R_{ij}\nabla_i v\nabla_j v + 2\nabla_i(\Delta v - |\nabla v|^2 + R - \frac{n}{2\tau})\nabla_i v \\ &= \Delta|\nabla v|^2 - 4R_{ij}\nabla_i v\nabla_j v - 2|\nabla^2 v|^2 - 2\nabla_i|\nabla v|^2\nabla_i v + 2\nabla_i R\nabla_i v,\end{aligned} \tag{3.3}$$

$$\begin{aligned}\frac{\partial\Delta v}{\partial\tau} &= -2R_{ij}\nabla_i\nabla_j v + \Delta v_\tau \\ &= \Delta^2 v - 2R_{ij}\nabla_i\nabla_j v - 2\nabla_i\Delta v\nabla_i v - 2|\nabla^2 v|^2 - 2R_{ij}\nabla_i v\nabla_j v + \Delta R,\end{aligned} \tag{3.4}$$

$$\frac{\partial R}{\partial\tau} = -\Delta R - 2|Rc|^2, \tag{3.5}$$

and

$$\frac{\partial}{\partial\tau}(\frac{v}{\tau}) = \Delta(\frac{v}{\tau}) - \frac{1}{\tau}|\nabla v|^2 + \frac{R}{\tau} - \frac{n}{2\tau^2} - \frac{v}{\tau^2}. \tag{3.6}$$

From the formulas above, the lemma follows immediately. □

By setting $b = -2$, $c = -1$ and $d = 0$, it follows from (3.2) that

$$\mathcal{L}F = -2\nabla_i F\nabla_i v - 2|R_{ij} + \nabla_i\nabla_j v|^2 + \frac{n}{\tau^2}.$$

Let $H = -|\nabla v|^2 + 2\Delta v + R$, then

$$\mathcal{L}H = -2\nabla_i H\nabla_i v - 2|R_{ij} + \nabla_i\nabla_j v|^2. \tag{3.7}$$

Moreover, we may write $H = 2z - y$, where $y = |\nabla v|^2 + R$ and $z = \Delta v + R$.

Proposition 3.2. *Let M^n be a complete manifold, and $g(t)$, $t \in [0,T]$, be a complete solution to the Ricci flow on M with bounded Ricci curvature and nonnegative scalar curvature. If u is a positive solution to (3.1), then for $v = -\ln u - \frac{n}{2}\ln(4\pi\tau)$, $\tau = T - t$, any point $O \in M$ and any constant $R_0 > 0$, we have*

$$-|\nabla v|^2 + 2\Delta v + R \le 2n[\frac{1}{\tau} + 16R_{max} + \frac{(72n+128)}{R_0^2} + \frac{8\sqrt{(n-1)R_{max}}}{R_0}]$$

on $\coprod_{\tau\in(0,T]} B_\tau(O,R_0)\times\{\tau\}$, where $R_{max} = \max_{[0,T]}|Rc|$.

Proof. Let $\phi(x,\tau) = \eta\left(\frac{r(x,\tau)}{2R_0}\right)$ on $\coprod_{\tau\in[0,T]} B_\tau(O,2R_0)\times\{\tau\}$, where $\eta(s)$ is the same function as in the proof of Proposition 2.3. From (3.7), we have

$$\tau\phi\mathcal{L}(\tau\phi H) = \tau\phi^2 H + \tau^2\phi H\phi_\tau - \tau^2\phi H\Delta\phi - 2\tau^2\phi\nabla_i\phi\nabla_i H - 2\tau^2\phi^2\nabla_i H\nabla_i v$$
$$- 2\tau^2\phi^2|R_{ij} + \nabla_i\nabla_j v|^2.$$

Let $\tilde{H} = \tau\phi H = 2\tilde{z} - \tilde{y}$, $\tilde{y} = \tau\phi y$, and $\tilde{z} = \tau\phi z$. Since

$$|\nabla\phi| \le \frac{8}{R_0}\phi^{\frac{1}{2}}, \quad \Delta\phi \ge -\frac{8n}{R_0^2} - \frac{8\sqrt{(n-1)R_{max}}}{R_0}, \quad \frac{\partial\phi}{\partial\tau} \le 16R_{max},$$

and

$$-2\tau_0^2\phi^2\nabla_i H\nabla_i v = 2\tau_0^2\phi H\nabla_i\phi\nabla_i v \le \frac{16}{R_0}\tau_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(2\tilde{z}-\tilde{y}),$$

if $\tilde{H}$ reaches a positive maximum at (x_0,τ_0), we have

$$\begin{aligned}
0 &\le \tau_0\phi\mathcal{L}(\tau\phi H)\\
&\le [1+16\tau_0 R_{max}+\frac{(8n+128)\tau_0}{R_0^2}+\frac{8\tau_0\sqrt{(n-1)R_{max}}}{R_0}](2\tilde{z}-\tilde{y})\\
&\quad+\frac{16}{R_0}\tau_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(2\tilde{z}-\tilde{y})-\frac{2}{n}|\tilde{z}|^2\\
&= [1+16\tau_0 R_{max}+\frac{(8n+128)\tau_0}{R_0^2}+\frac{8\tau_0\sqrt{(n-1)R_{max}}}{R_0}](2\tilde{z}-\tilde{y})\\
&\quad+\frac{16}{R_0}\tau_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(2\tilde{z}-\tilde{y})-\frac{1}{2n}(2\tilde{z}-\tilde{y}+\tilde{y})^2\\
&\le -\frac{1}{2n}(2\tilde{z}-\tilde{y})^2+[1+16\tau_0 R_{max}+\frac{(72n+128)\tau_0}{R_0^2}+\frac{8\tau_0\sqrt{(n-1)R_{max}}}{R_0}](2\tilde{z}-\tilde{y}).
\end{aligned}$$

Therefore, at (x_0,τ_0), we have

$$\tilde{H}\le 2n[1+16\tau_0 R_{max}+\frac{(72n+128)\tau_0}{R_0^2}+\frac{8\tau_0\sqrt{(n-1)R_{max}}}{R_0}].$$

Hence for any (x,τ), one has

$$\tau H\le 2n[1+16\tau R_{max}+\frac{(72n+128)\tau}{R_0^2}+\frac{8\tau\sqrt{(n-1)R_{max}}}{R_0}].$$

□

Letting $R_0\to\infty$ in the proposition above, we have

Corollary 3.3. *Let M^n be a complete manifold, and $g(t)$, $t\in[0,T]$, be a complete solution to the Ricci flow on M with bounded Ricci curvature and nonnegative scalar curvature. If u is a positive solution to* (3.1)*, then for $v=-\ln u-\frac{n}{2}\ln(4\pi\tau)$, $\tau=T-t$, we have*

$$-|\nabla v|^2+2\Delta v+R\le 2n[\frac{1}{\tau}+16R_{max}],$$

on $M\times[0,T)$, where $R_{max}=\max_{M\times[0,T]}|Rc|$.

Now we are ready to proveTheorem 1.3.

Proof of Theorem 1.3: Let $\hat{H}=\tau H-k$, then (3.7) implies that

$$\tau\mathcal{L}\hat{H}=\hat{H}+k-2\tau\nabla_i\hat{H}\nabla_i v-2\tau^2|R_{ij}+\nabla_i\nabla_j v|^2.$$

Let $\check{H} = \hat{H} - \epsilon\psi$, where $\psi = e^{A\tau} f$ for some constant A to be determined and f is the same function as in Lemma 2.5, then

$$\begin{aligned}\tau\mathcal{L}(\check{H}) &= \tau\mathcal{L}(\hat{H}) - A\tau\epsilon\psi + \tau\epsilon e^{A\tau}\Delta f \\ &\le \check{H} + k + \epsilon\psi - 2\tau\nabla_i\check{H}\nabla_i v - 2\tau\epsilon e^{A\tau}\nabla_i f\nabla_i v - \frac{1}{2n}(\check{H} + k + \epsilon\psi + \tau y)^2 - (A-C)\tau\epsilon\psi \\ &\le \check{H} + k + \epsilon\psi - 2\tau\nabla_i\check{H}\nabla_i v + n\tau\epsilon e^{A\tau}|\nabla f|^2 + \frac{1}{n}\tau\epsilon e^{A\tau}|\nabla v|^2 \\ &\quad - \frac{1}{2n}(\check{H} + k + \epsilon\psi + \tau y)^2 - (A-C)\tau\epsilon\psi.\end{aligned}$$

Assume that at time τ_0 and some point x_0, $\tilde{H}$ reaches 0 for the first time, then

$$\begin{aligned}0 \le \tau_0\mathcal{L}(\check{H}) &\le k + \epsilon\psi + n\tau_0\epsilon e^{A\tau_0}|\nabla f|^2 + \frac{1}{n}\tau_0\epsilon e^{A\tau_0}|\nabla v|^2 - \frac{1}{2n}(k + \epsilon\psi + \tau_0 y)^2 - (A-C)\tau_0\epsilon\psi \\ &\le k - \frac{k^2}{2n} + \epsilon\psi - \frac{k}{n}\epsilon\psi + \frac{1}{n}\tau_0\epsilon\psi y - \frac{1}{n}(k+\epsilon\psi)\tau_0 y - \frac{1}{2n}\tau_0^2 y^2 - [A - (n+1)C]\tau_0\epsilon\psi.\end{aligned}$$

Thus, if we pick $k = 2n$ and $A > (n+1)C$, there is a contradiction. It means that

$$\hat{H} - \epsilon\psi < 0$$

for any $\epsilon > 0$.

Letting $\epsilon \to 0$ yields $\hat{H} \le 0$, i.e.,

$$2\Delta v - |\nabla v|^2 + R \le \frac{2n}{\tau}.$$

□

Remark 3.4. *Theorem 1.3 generalizes the Li-Yau type estimate* (1.8) *obtained by X. Cao* [9] *and Kuang-Zhang* [23] *to the complete noncompact setting.*

4. Li-Yau-Hamilton estimates for the heat equation

In this last section, we apply our arguments in Section 2 to the heat equation with potential under the Ricci flow and derive an Li-Yau type estimate for positive solutions on complete noncompact manifolds, extening the previous work of X. Cao and R. Hamilton [10] in the compact case. More precisely, we consider the following heat equation with a linear forcing term,

$$\frac{\partial u}{\partial t} = \Delta u + Ru, \tag{4.1}$$

coupled with the Ricci flow (1.5).

For any positive solution u to (4.1), let $v = -\ln u$. Then, it is easy to check that

$$\frac{\partial v}{\partial t} = \Delta v - |\nabla v|^2 - R.$$

Let

$$\begin{aligned}F &= |\nabla v|^2 + bv_t + cR \\ &= b\Delta v + (1-b)|\nabla v|^2 + (c-b)R\end{aligned}$$

and

$$\mathcal{L} = \frac{\partial}{\partial t} - \Delta.$$

Similarly to Proposition 2.1 in [7], one can show

Proposition 4.1.

$$\mathcal{L}(F) = -2\nabla_i F\nabla_i v - 2|\nabla^2 v|^2 + 2bR_{ij}\nabla_i\nabla_j v + 2c|Rc|^2 \\ - b\frac{\partial R}{\partial t} + 2(c-1)\nabla_i R\nabla_i v - 2bR_{ij}\nabla_i v\nabla_j v.$$

By choosing appropriate c and a cut-off function, it is not hard to drive the following local differential Harnack estimate.

Proposition 4.2. *Let $(M^n, g_{ij}(t))$, $t \in [0,T]$, be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator. If u is a positive solution to (4.1), then for $v = -\ln u$, $b > 1$, any point $O \in M$ and any constant $R_0 > 0$, we have*

$$|\nabla v|^2 + bv_t + (1-b)R \le \frac{b^2 n}{2}\left[\frac{1}{t} + 16R_{max} + \frac{C_0}{R_0^2} + C_1(R_{max}, t)\right] \tag{4.2}$$

on $\coprod_{t\in(0,T]} B_t(O,R_0)\times\{t\}$ when $1 < b \le 2$, and

$$|\nabla v|^2 + bv_t + (1-b)R \le \frac{b^2 n}{2}\left[\frac{1}{t} + (16 + \frac{2(b-2)}{bn})R_{max} + \frac{C_0}{R_0^2} + C_1(R_{max}, t)\right] \tag{4.3}$$

on $\coprod_{t\in(0,T]} B_t(O,R_0)\times\{t\}$ when $b \ge 2$. Here $C_0 = 8n + 128 + \frac{16b^2 n}{b-1}$, $C_1(R_{max}, t) = \sqrt{\frac{2}{b^2 n}[\frac{(b-2)^2}{2}R_{max}^2 + \frac{bR_{max}}{t}]}$, and $R_{max} = \sup_{M\times[0,T]} R$.

Proof. By taking $c = 1 - b$, we have

$$F = |\nabla v|^2 + bv_t + (1-b)R = y + bz,$$

where $y = |\nabla v|^2 + R$ and $z = v_t - R$. Hence, by Proposition 4.1,

$$\mathcal{L}(F) = -2\nabla_i F\nabla_i v - 2|\nabla^2 v - \frac{b}{2}Rc|^2 + \frac{(b-2)^2}{2}|Rc|^2 \tag{4.4}$$

$$- b[R_t + 2\nabla_i R\nabla_i v + 2R_{ij}\nabla_i v\nabla_j v]. \tag{4.5}$$

Let $\phi(x,t)$ be the cut-off function in (2.3). Then, it follows that

$$t\phi\mathcal{L}(t\phi F) = \phi(\tilde{y} + b\tilde{z}) + t\phi_t(\tilde{y} + b\tilde{z}) - t\Delta\phi(\tilde{y} + b\tilde{z}) - 2t^2\phi\nabla_i\phi\nabla_i F - 2t^2\phi^2\nabla_i F\nabla_i v \\ - 2t^2\phi^2|\nabla^2 v - \frac{b}{2}Rc|^2 + \frac{(b-2)^2}{2}t^2\phi^2|Rc|^2 - bt\phi^2 Q + bt\phi^2 R,$$

where Q is the quantity in (2.4), $\tilde{y} = t\phi y$, and $\tilde{z} = t\phi z$.

Recall that

$$|\frac{\partial\phi}{\partial t}| \le 16R_{max}, \qquad \Delta\phi \ge -\frac{8n}{R_0^2}, \qquad |\nabla\phi| \le \frac{8}{R_0}\phi^{\frac{1}{2}}.$$

If $t\phi F \le 0$ in $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, then we are done. Otherwise, since $t\phi F = 0$ on the parabolic boundary of $\coprod_{t\in[0,T]} B_t(O,2R_0)\times\{t\}$, we may assume that $t\phi F$ achieves a positive maximum for the first time at some $t_0 > 0$ and some interior point x_0. Thus at (x_0,t_0), we have

$$\tilde{y} + b\tilde{z} = t_0\phi F > 0, \qquad F\nabla\phi = -\phi\nabla F, \qquad \text{and} \qquad \mathcal{L}(t\phi F)(x_0,t_0) \ge 0.$$

Since

$$-2t_0^2\phi^2\nabla_i F\nabla_i v = 2t_0^2\phi F\nabla_i\phi\nabla_i v \le \frac{16}{R_0}t_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(\tilde{y} + b\tilde{z}),$$

we have

$$\begin{aligned}
0 &\le t_0\phi\mathcal{L}(t\phi F)\\
&\le (\tilde{y}+b\tilde{z})+16t_0R_{max}(\tilde{y}+b\tilde{z})+\frac{(8n+128)t_0}{R_0^2}(\tilde{y}+b\tilde{z})+\frac{16}{R_0}t_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(\tilde{y}+b\tilde{z})\\
&\quad-\frac{2}{n}t_0^2\phi^2(y+z+\frac{2-b}{2}R)^2+\frac{(b-2)^2}{2}t_0^2\phi^2R^2+bt_0\phi^2R\\
&\le (\tilde{y}+b\tilde{z})\left[1+16t_0R_{max}-\frac{4(b-1)}{b^2n}\tilde{y}+\frac{16}{R_0}t_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}+\frac{(8n+128)t_0}{R_0^2}\right]-\frac{2(b-1)^2}{b^2n}\tilde{y}^2\\
&\quad-\frac{2}{b^2n}(\tilde{y}+b\tilde{z})^2+\frac{(b-2)^2(n-1)}{2n}t_0^2\phi^2R^2-\frac{2(b-1)(2-b)}{bn}\tilde{y}t_0\phi R\\
&\quad-\frac{2(2-b)}{bn}t_0\phi R(\tilde{y}+b\tilde{z})+bt_0\phi R.
\end{aligned}$$

If $1<b\le 2$, we have

$$\begin{aligned}
0 &\le -\frac{2}{b^2n}(\tilde{y}+b\tilde{z})^2+\left[1+16t_0R_{max}-\frac{4(b-1)}{b^2n}\tilde{y}+\frac{16}{R_0}t_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}+\frac{(8n+128)t_0}{R_0^2}\right](\tilde{y}+b\tilde{z})\\
&\quad+\frac{(b-2)^2(n-1)}{2n}t_0^2R_{max}^2+bt_0R_{max}\\
&\le -\frac{2}{b^2n}(\tilde{y}+b\tilde{z})^2+\left[1+16t_0R_{max}+(8n+128+\frac{16b^2n}{b-1})\frac{t_0}{R_0^2}\right](\tilde{y}+b\tilde{z})\\
&\quad+\frac{(b-2)^2(n-1)}{2n}t_0^2R_{max}^2+bt_0R_{max},
\end{aligned}$$

which implies (4.2).

If $b\ge 2$, then

$$\begin{aligned}
0 &\le (\tilde{y}+b\tilde{z})+16t_0R_{max}(\tilde{y}+b\tilde{z})+\frac{(8n+128)t_0}{R_0^2}(\tilde{y}+b\tilde{z})+\frac{16}{R_0}t_0^{\frac{1}{2}}\tilde{y}^{\frac{1}{2}}(\tilde{y}+b\tilde{z})-\frac{2(b-1)^2}{b^2n}\tilde{y}^2\\
&\quad-\frac{2}{b^2n}(\tilde{y}+b\tilde{z})^2+\frac{(b-2)^2(n-1)}{2n}t_0^2\phi^2R^2-\frac{4(b-1)}{b^2n}\tilde{y}(\tilde{y}+b\tilde{z})+\frac{2(b-1)(b-2)}{bn}\tilde{y}t_0\phi R\\
&\quad+\frac{2(b-2)}{bn}t_0\phi R(\tilde{y}+b\tilde{z})+bt_0\phi R\\
&\le -\frac{2}{b^2n}(\tilde{y}+b\tilde{z})^2+\left[1+(16+\frac{2(b-2)}{bn})t_0R_{max}+(8n+128+\frac{16b^2n}{b-1})\frac{t_0}{R_0^2}\right](\tilde{y}+b\tilde{z})\\
&\quad+\frac{(b-2)^2}{2}t_0^2R_{max}^2+bt_0R_{max},
\end{aligned}$$

which leads to (4.3). This finishes the proof. □

Letting $R_0\to\infty$, we have

Corollary 4.3. *Let $(M^n,g_{ij}(t))$, $t\in[0,T]$, be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator. If u is a positive solution to (4.1), then for $v=-\ln u$, we have*

$$|\nabla v|^2+bv_t+(1-b)R\le\frac{b^2n}{2}\left[\frac{1}{t}+16R_{max}+C_1(R_{max},t)\right]$$

on $M\times(0,T]$ when $1<b\le 2$, and

$$|\nabla v|^2+bv_t+(1-b)R\le\frac{b^2n}{2}\left[\frac{1}{t}+(16+\frac{2(b-2)}{bn})R_{max}+C_1(R_{max},t)\right]$$

on $M\times(0,T]$ when $b\ge 2$. Here R_{max} and $C_1(R_{max},t)$ are the same constants as in Proposition 4.2.

Again, the above result can be refined as

Theorem 4.4. *Let $(M^n,g_{ij}(t))$, $t\in[0,T]$, be a complete solution to the Ricci flow with bounded curvature and nonnegative curvature operator. If u is a positive solution to (4.1), then for $v=-\ln u$, we have*

$$|\nabla v|^2+bv_t+(1-b)R-\frac{b(2-b)\sqrt{n-1}}{2}R_{max}-\frac{b^2n}{2t}\le 0$$

on $M\times(0,T]$ when $1<b\le 2$, and

$$|\nabla v|^2+bv_t+(1-b)R-b(b-2)[1+\frac{\sqrt{n-1}}{2}]R_{max}-\frac{b^2n}{2t}\le 0$$

on $M\times(0,T]$ when $b\ge 2$. Here $R_{max}=\sup_{M\times[0,T]}R$.

Proof. Let $H=t(F-K)-d$ with d and K to be determined, from (4.4) one has

$$\begin{aligned}t\mathcal{L}(H)\le&H+d-2t\nabla_iH\nabla_iv-\frac{2}{b^2n}\left[H+tK+d+(b-1)ty+\frac{b(2-b)}{2}tR\right]^2\\&+\frac{(b-2)^2}{2}t^2R^2+btR\\=&H+d-2t\nabla_iH\nabla_iv-\frac{2}{b^2n}(H+tK+d)^2-\frac{2(b-1)^2}{b^2n}t^2y^2+\frac{(b-2)^2(n-1)}{2n}t^2R^2\\&-\frac{4(b-1)}{b^2n}ty(H+tK+d)-\frac{2(2-b)}{bn}tR(H+tK+d)-\frac{2(b-1)(2-b)}{bn}t^2Ry+btR.\end{aligned}$$

Let $\tilde H=H-\epsilon\psi$, where $\psi=e^{At}f$ with some constant A to be determined and f being the same function as in Lemma 2.5. From Corollary 4.3, we know that $\tilde H<0$ at $t=0$ and outside a fixed compact subset of M for $t\in(0,T]$. Suppose that $\tilde H=0$ for the first time at $t=t_0>0$ and some point $x_0\in M$. Then, at (x_0,t_0), we have

$$H=\epsilon\psi>0,\qquad\nabla\tilde H=0,\qquad\text{and}\qquad\mathcal{L}(\tilde H)\ge 0.$$

If $1<b\le 2$ and choose $d=\frac{b^2n}{2}$, then

$$\begin{aligned}0\le&\,t_0\mathcal{L}(\tilde H)\\\le&\,\epsilon\psi+d+2t_0\epsilon e^{At_0}\nabla_if\nabla_iv-\frac{2}{b^2n}(\epsilon\psi+t_0K+d)^2-\frac{2(b-1)^2}{b^2n}t_0^2y^2+\frac{(b-2)^2(n-1)}{2n}t_0^2R^2\\&-\frac{4(b-1)}{b^2n}t_0y(\epsilon\psi+t_0K+d)-\frac{2(2-b)}{bn}t_0R(\epsilon\psi+t_0K+d)-\frac{2(b-1)(2-b)}{bn}t_0^2Ry\\&+bt_0R-At_0\epsilon\psi+t_0\epsilon e^{At_0}\Delta f\\\le&-\epsilon\psi-\frac{2}{b^2n}(\epsilon\psi)^2+\delta t_0\epsilon\psi|\nabla v|^2+\frac{C}{4\delta}t_0\epsilon\psi-\frac{2(b-1)^2}{b^2n}t_0^2|\nabla v|^4\\&+\frac{(b-2)^2(n-1)}{2n}t_0^2R_{max}^2-\frac{2}{b^2n}t_0^2K^2-(2-b)(b-1)t_0R-At_0\epsilon\psi+Ct_0\epsilon\psi.\end{aligned}$$

Now if we choose $K = \frac{b(2-b)\sqrt{n-1}}{2} R_{max}$, and some $\delta > 0$ such that

$$-\frac{2}{b^2 n}(\epsilon\psi)^2 + \delta t_0 \epsilon\psi |\nabla v|^2 - \frac{2(b-1)^2}{b^2 n} t_0^2 |\nabla v|^4 \leq 0,$$

then we deduce

$$0 \leq [(1 + \frac{1}{4\delta})C - A] t_0 \epsilon\psi.$$

By further choosing $A > (1 + \frac{1}{4\delta})C$, we get a contradiction.

If $b \geq 2$ and $d = \frac{b^2 n}{2}$, then

$$\begin{aligned}
0 &\leq t_0 \mathcal{L}(\tilde{H}) \\
&\leq \epsilon\psi + d + 2t_0 \epsilon e^{At_0} \nabla_i f \nabla_i v - \frac{2}{b^2 n}(\epsilon\psi + t_0 K + d)^2 - \frac{2(b-1)^2}{b^2 n} t_0^2 y^2 + \frac{(b-2)^2 (n-1)}{2n} t_0^2 R^2 \\
&\quad - \frac{4(b-1)}{b^2 n} t_0 y (\epsilon\psi + t_0 K + d) + \frac{2(b-2)}{bn} t_0 R (\epsilon\psi + t_0 K + d) + \frac{2(b-1)(b-2)}{bn} t_0^2 R y \\
&\quad + b t_0 R - A t_0 \epsilon\psi + t_0 \epsilon e^{At_0} \Delta f \\
&\leq -\epsilon\psi - (\frac{4}{b^2 n} K - \frac{2(b-2)}{bn} R) t_0 \epsilon\psi - \frac{2}{b^2 n}(\epsilon\psi)^2 + \delta t \epsilon\psi |\nabla v|^2 + \frac{C}{4\delta} t_0 \epsilon\psi - \frac{2(b-1)^2}{b^2 n} t_0^2 |\nabla v|^4 \\
&\quad + \frac{(b-2)^2 (n-1)}{2n} t_0^2 R^2 + \frac{2(b-2)}{bn} t_0^2 R K - \frac{2}{b^2 n} t_0^2 K^2 + (b-2)(b-1) t_0 R - 2 t_0 K \\
&\quad - A t_0 \epsilon\psi + C t_0 \epsilon\psi - \frac{2(b-1)}{bn} [\frac{2}{b} K - (b-2) R] t_0^2 y.
\end{aligned}$$

Let $K = b(b-2)(1 + \frac{\sqrt{n-1}}{2}) R_{max}$, then

$$\frac{2}{b} K - (b-2) R \geq 0,$$

$$(b-2)(b-1) R - 2K \leq 0,$$

and

$$\frac{(b-2)^2 (n-1)}{2n} R^2 + \frac{2(b-2)}{bn} R K - \frac{2}{b^2 n} K^2 \leq 0.$$

If we further choose some appropriate $\delta > 0$ so that

$$-\frac{2}{b^2 n}(\epsilon\psi)^2 + \delta t \epsilon\psi |\nabla v|^2 - \frac{2(b-1)^2}{b^2 n} t_0^2 |\nabla v|^4 \leq 0,$$

then

$$0 \leq t_0 \mathcal{L}(\tilde{H}) \leq (-A + C + \frac{C}{4\delta}) t_0 \epsilon\psi.$$

By choosing $A > C + \frac{C}{4\delta}$, we also get a contradiction.

Therefore, in either case, it must be true that $\tilde{H} < 0$. Letting $\epsilon \to 0$, we conclude that $H \leq 0$. This finishes the proof of Theorem 3.4. □

In the special case when $b = 2$, we obtain Theorem 1.4.

Remark 4.5. *When M is compact, Theorem 1.4 was first proved by X. Cao and R. Hamilton in* [10].

References

[1] Aronson, D. G., and Bénilan, P. Régularité des solutions de l'équation des milieux poreux dans $\mathbf{R}^N$. *C. R. Acad. Sci. Paris Sér. A-B 288*, 2 (1979), A103–A105.

[2] Bailesteanu, M., Cao, X., and Pulemotov, A. Gradient estimates for the heat equation under the Ricci flow. *J. Funct. Anal. 258*, 10 (2010), 3517–3542.

[3] Bernard, G. Existence theorems for fast diffusion equations. *Nonlinear Anal. 43*, 5, Ser. A: Theory Methods (2001), 575–590.

[4] Brendle, S. A generalization of Hamilton's differential Harnack inequality for the Ricci flow. *J. Differential Geom. 82*, 1 (2009), 207–227.

[5] Cao, H. D. On Harnack's inequalities for the Kähler-Ricci flow. *Invent. Math. 109*, 2 (1992), 247–263.

[6] Cao, H.-D., and Ni, L. Matrix Li-Yau-Hamilton estimates for the heat equation on Kähler manifolds. *Math. Ann. 331*, 4 (2005), 795–807.

[7] Cao, H.-D., and Zhu, M. Aronson-Bénilan estimates for the porous medium equation under the Ricci flow. *J. Math. Pures Appl. (9) 104*, 4 (2015), 729–748.

[8] Cao, H.-D., and Zhu, X.-P. A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow. *Asian J. Math. 10*, 2 (2006), 165–492.

[9] Cao, X. Differential Harnack estimates for backward heat equations with potentials under the Ricci flow. *J. Funct. Anal. 255*, 4 (2008), 1024–1038.

[10] Cao, X., and Hamilton, R. S. Differential Harnack estimates for time-dependent heat equations with potentials. *Geom. Funct. Anal. 19*, 4 (2009), 989–1000.

[11] Chow, B. The Ricci flow on the 2-sphere. *J. Differential Geom. 33*, 2 (1991), 325–334.

[12] Dahlberg, B. E. J., and Kenig, C. E. Nonnegative solutions to fast diffusions. *Rev. Mat. Iberoamericana 4*, 1 (1988), 11–29.

[13] Daskalopoulos, P., and Del Pino, M. On nonlinear parabolic equations of very fast diffusion. *Arch. Rational Mech. Anal. 137*, 4 (1997), 363–380.

[14] Daskalopoulos, P., and del Pino, M. On the Cauchy problem for $u_t = \Delta \log u$ in higher dimensions. *Math. Ann. 313*, 2 (1999), 189–206.

[15] Daskalopoulos, P., and del Pino, M. A. On fast diffusion nonlinear heat equations and a related singular elliptic problem. *Indiana Univ. Math. J. 43*, 2 (1994), 703–728.

[16] Daskalopoulos, P., and del Pino, M. A. On a singular diffusion equation. *Comm. Anal. Geom. 3*, 3-4 (1995), 523–542.

[17] Daskalopoulos, P., Hamilton, R., and Sesum, N. Classification of ancient compact solutions to the Ricci flow on surfaces. *J. Differential Geom. 91*, 2 (2012), 171–214.

[18] Hamilton, R. S. The Ricci flow on surfaces. In *Mathematics and general relativity (Santa Cruz, CA, 1986)*, vol. 71 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 1988, pp. 237–262.

[19] Hamilton, R. S. The Harnack estimate for the Ricci flow. *J. Differential Geom. 37*, 1 (1993), 225–243.

[20] Hamilton, R. S. A matrix Harnack estimate for the heat equation. *Comm. Anal. Geom. 1*, 1 (1993), 113–126.

[21] Herrero, M. A., and Pierre, M. The Cauchy problem for $u_t = \Delta u^m$ when $0 < m < 1$. *Trans. Amer. Math. Soc. 291*, 1 (1985), 145–158.

[22] King, J. R. Interacting dopant diffusions in crystalline silicon. *SIAM J. Appl. Math. 48*, 2 (1988), 405–415.

[23] Kuang, S., and Zhang, Q. S. A gradient estimate for all positive solutions of the conjugate heat equation under Ricci flow. *J. Funct. Anal. 255*, 4 (2008), 1008–1023.

[24] Li, P., and Yau, S.-T. On the parabolic kernel of the Schrödinger operator. *Acta Math. 156*, 3-4 (1986), 153–201.

[25] Lions, P. L., and Toscani, G. Diffusive limit for finite velocity Boltzmann kinetic models. *Rev. Mat. Iberoamericana 13*, 3 (1997), 473–513.

[26] Lu, P., Ni, L., Vázquez, J.-L., and Villani, C. Local Aronson-Bénilan estimates and entropy formulae for porous medium and fast diffusion equations on manifolds. *J. Math. Pures Appl. (9) 91*, 1 (2009), 1–19.

[27] Ni, L. A note on Perelman's LYH-type inequality. *Comm. Anal. Geom. 14*, 5 (2006), 883–905.

[28] Ni, L. A matrix Li-Yau-Hamilton estimate for Kähler-Ricci flow. *J. Differential Geom. 75*, 2 (2007), 303–358.

[29] Okuda, H., and Dawson, J. M. Theory and numerical simulation on lasma diffusion accross a magnetic field. *Phys. Fulids 16* (1973), 408–426.

[30] Perelman, G. The entropy formula for the ricci flow and its geometric applications. *arXiv:math/0211159* (2002).

[31] Vázquez, J. L. Nonexistence of solutions for nonlinear heat equations of fast-diffusion type. *J. Math. Pures Appl. (9) 71*, 6 (1992), 503–526.
[32] Vázquez, J. L. *Smoothing and decay estimates for nonlinear diffusion equations*, vol. 33 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2006. Equations of porous medium type.
[33] Vázquez, J. L. *The porous medium equation*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2007. Mathematical theory.
[34] Wu, L.-F. A new result for the porous medium equation derived from the Ricci flow. *Bull. Amer. Math. Soc. (N.S.) 28*, 1 (1993), 90–94.
[35] Zhang, Q. S., and Zhu, M. Li-yau gradient bounds under nearly optimal curvature conditions. *arXiv:1511.00791* (2015).

Huai-Dong Cao, Department of Mathematics, Lehigh University, Bethlehem, PA 18015, USA
E-mail address: huc2@lehigh.edu

Meng Zhu, Department of Mathematics, East China Normal University, Shanghai 200241, China & Department of Mathematics, University of California at Riverside, Riverside, CA 92521, USA
E-mail address: mzhu@math.ecnu.edu.cn, mengzhu@ucr.edu